

Diverse Noachian Environments

Nili Fossae Trough

*Jack Mustard, Bethany Ehlmann, J. R. Skok,
Dave Des Marais, Nicolas Mangold, and Francois Poulet*

Investigation of Diverse Habitable Noachian Environments
(Hydrothermal, alluvial/fluvial, shallow crust/pedogenic)

Sampling and characterizing

Impact ejecta

Hesperian volcanics

Phyllosilicate-bearing infill of Nili Fossae

Strongly altered Noachian crust

Unaltered Noachian crust

Fully Address E2E SAG Top 3 Objectives

Life (A1) Planetary Evolution (B1)

Water (C1)



Noachian crust enriched in low-
Ca pyroxene




Noachian crust enriched in
phyllosilicate

NOTE ADDED BY JPL WEBMASTER: This content has not been approved or adopted by NASA, JPL, or the California Institute of Technology. This document is being made available for information purposes only, and any views and opinions expressed herein do not necessarily state or reflect those of NASA, JPL, or the California Institute of Technology.


Multiple, Distinct Environments

- Noachian is when phyllosilicate formation was most intense: Access to the source environments
- Hydrothermal systems
 - Fractures as transport pathways for water, mineralization in fractures
 - No mounds, spring deposits, but those are surface features and not expected to survive to the present
- Sedimentary units
 - Infill of trough, formation of sapping channel
 - Regionally, layered units in crater floors, troughs
 - Erosion in go-to site leaves outcrops of remnants of these processes
- Subsurface groundwater or shallow crustal environment
 - Protected from destructive radiation environment
 - Abundant chemical energy sources
 - Fluid flow in the crust/groundwater transported nutrients/energy




Broad Mars Scientific Objectives

- Noachian Habitable Environments
 - Ancient crustal and genesis region
 - Fluvially transported sediments
 - Hydrothermal systems
- Impact processes
 - Superbly exposed ejecta from 65 km Hargraves crater
 - Ejecta blocks in a phyllosilicate-bearing matrix
 - Transport, fluidization, alteration
- Composition and character of ancient, unaltered crust
- Composition, mineralogy, and texture of Hesperian Syrtis Major lava: A datable surface the time stratigraphic marker
- Traverse the Noachian-Hesperian Boundary
- Phyllosilicate-Transported
- Phyllosilicate-In Place



Stratigraphy and Processes in Isidis-Nili Fossae



Isidis Basin and Syrtis
Major lavas are major time-
stratigraphic markers


Significant gradation
(sedimentary? aeolian?
alluvial?) between Isidis
basin formation and Syrtis
lava emplacement





Clearly defined wet periods

Nili Fossae Trough

- Diverse Noachian environments present throughout the landing site
- Regional geology, represented in the landing site, indicates sustained interaction of water with the crust over an extended period as a consequence of multiple episodes of distinct character
 - Fe/Mg Phyllosilicates with variation in band position, strength of water absorption
 - Smectite clay transported and deposited in fluvial systems
 - Regional episode of kaolinite formation
 - Carbonate formation in association with olivine
 - Chlorite, zeolite, and hydrated silicate in association with impacts
- The region north east of Syrtis Major was persistently wet and the geologic context for understanding the interaction of water is extraordinarily well preserved and exposed
- The Nili Fossae Trough landing site sits within this region and provides exciting access to a diverse suite of environments

- Olivine
- Low-Ca Pyroxene
- Phyllosilicate
- Fe-Phyllosilicate





-  Olivine
-  Low-Ca Pyroxene
-  Phyllosilicate
-  Fe-Phyllosilicate

CRISM Observations

FRT00007BC8

FRT000064D9






Unaltered basement enriched
in low-Ca pyroxene as capping
unit and on the floor



Thick section of phyllosilicate-
rich Noachian basement
exposed in Fossae walls

Phyllosilicate-bearing
basement beneath volcanics

Syrtis Major Volcanics




FRT000064D9:
2.4, 1.8, 1.15 μm RGB








Mineralogy identified


Fe-oxide and crystalline hematite
Fe/Mg Smectite with variety of
band positions, H₂O content
Kaolinite
Carbonate
Pyroxene (Low and High Ca)
Olivine

- Scarp walls
- Noachian crust
- Channel floor
- Trough fill
- Hesperian lava
- Hargraves Ejecta
- Fresh Craters





-  Olivine
-  Low-Ca Pyroxene
-  Phyllosilicate
-  Fe-Phyllosilicate









(Representative vertical and horizontal distances, not to scale)




- Low-Ca Pyroxene
- Phyllosilicate
- Fe-Phyllosilicate




5 km




250 m





(Representative vertical and horizontal distances, not to scale)

Widespread gradation by alluvial
and fluvial processes has filled in
topographic lows
1 km thickness in impact craters
Deposited in Nili Fossae troughs





Layered material on the
floor of the sapping
channel


Trough fill

Transported phyllosilicate-
bearing alluvial/fluvial
deposits in the go-to site


Phyllosilicate-bearing sediments
1.91 and 2.29 μm absorptions











5x vertical exaggeration




(Representative vertical and horizontal distances, not to scale)







- Low-Ca Pyroxene
- Phyllosilicate
- Fe-Phyllosilicate



500 m

Unaltered Noachian
crust embedded in
phyllosilicate-bearing
formation

HiRISE Color

Layered
Sediments



Fe/Mg smectite

Low-Ca pyroxene



75 m

27

Fractures, conduits for flow, mineralization



- Low-Ca Pyroxene
- Phyllosilicate
- Fe-Phyllosilicate



Nili Fossae Trough

- Diverse habitable environments mitigates against a limited or pre-judged suite of environments
- Broader science goals advance understanding of Mars as a planet
- Diverse Noachian environments present throughout the landing site
- Regional geologic context, represented in the landing site, indicates sustained interaction of water with the crust as a consequence of multiple episodes of distinct character
 - Fe/Mg Phyllosilicates with variation in band position, strength of water absorption
 - Smectite clay transported and deposited in fluvial systems
 - Regional episode of kaolinite formation
 - Carbonate formation in association with olivine
- The region north east of Syrtis Major was persistently wet and the geologic context for understanding the interaction of water is extraordinarily well preserved and exposed